

Teen Abortion Risks Fact Sheet

EXHIBIT 3
DATE 3/23/2011
SB 97

"Parents are faced with a shell of a person and have no idea where they lost their child."
—Terri, who had a secret abortion as a teen

Suicide attempts 6 times more likely

- **Teenagers are 6 times more likely to attempt suicide if they have had an abortion in the last six months** than are teens who have not had an abortion,¹ and four times more likely to commit suicide than adults who abort.² A history of abortion is likely to be associated with adolescent suicidal thinking.¹ Overall, women who have abortions have a 6 times higher rate of suicide compared to women who carry to term.³
- **Teens who abort are more likely to develop psychological problems,**⁴ and are nearly three times more likely to be admitted to mental health hospitals than teens in general.⁵
- **About 40% of teen abortions take place with no parental involvement,**⁶ leaving parents in the dark about subsequent emotional or physical problems and putting them at risk for further injury and death.⁷
- **Teens are 5 times more likely to seek subsequent help for psychological and emotional problems** compared to their peers who carry "unwanted pregnancies" to term.⁸
- **Teens are 3 times more likely to report subsequent trouble sleeping, and nine times more likely to report subsequent marijuana use after abortion.**⁸
- **Among studies comparing abortion vs. carrying to term, worse outcomes are associated with abortion,** even when the pregnancy is unplanned.⁸
- **Teens are more likely to abort because of pressure from their parents or partner.**⁹ In one study, 64% of women who had undergone an abortion reported that they felt pressured by others to abort.¹⁰
- **Teens are more likely to report being misinformed in pre-abortion counseling.**¹¹

Acute Pain. Infertility. Risk of Death.

- **Acute pain.** Teens report more severe pain during the abortion procedure than do adult women.¹²
- **Lacerations up to twice as likely.** Teens are up to twice as likely to experience dangerous cervical lacerations during abortion compared to older women.¹³
- **Infertility and life-threatening complications.** Studies have found that teens are at higher risk for post-abortion infections such as pelvic inflammatory disease (PID) and endometritis.¹⁴ These infections increase their risk of infertility, hysterectomy, ectopic pregnancy, and other serious complications.¹⁵
- **Higher breast cancer risk.** An early full-term birth reduces breast cancer risk by as much as 1/3, while abortion of a first pregnancy carries a 30- to 50% increased risk of breast cancer.¹⁶ More than 90% of those who abort at 17 or younger have not had a previous full-term pregnancy, compared to 78% of patients age 18-19 and 49% of abortion patients overall.¹⁷
- **Teens are more likely to have riskier late-term abortions.** According to the CDC, approximately 30% of abortions among teens take place at 13 weeks gestation or greater, compared to only 12% among women in general.¹⁸ Late-term abortions are associated with more severe psychological problems,¹⁹ higher risk of physical complications,²⁰ and problems in later pregnancies.²¹

Grief, trauma and self-destructive outcomes

- **Teens who abort are twice as likely as their peers to abuse alcohol, marijuana, or cocaine.**²²

continued ▶

- * **Teens have greater difficulty coping** after abortion,²³ leading to problems such as suicide, psychological problems, substance abuse, and difficulty in relationships.
- **Negative effects on relationships and parenting.** Teens who report “being particularly fond of children” do not do as well psychologically after an abortion.²⁴ Teens who have abortions often have later problems regarding sexuality and parenting.²⁵
- **A lonely, traumatic experience.** The abortion procedure itself is considered by many teens to be stressful and associated with feelings of guilt, depression, and a sense of isolation.²⁶
- **A nightmare that doesn't end.** Teens are more likely to report severe nightmares and to score higher on scales measuring antisocial traits, paranoia, drug abuse, and psychotic delusions than are older abortion patients.²⁷
- **Four times higher risk of repeat abortion.** Teens who abort are likely to become pregnant again within the next few years.²⁸ Among pregnant teens, those who had had an abortion were at least 4 times more likely to abort.²⁹

Citations

1. Garfinkel et al., “Stress, Depression and Suicide: A Study of Adolescents in Minnesota,” *Responding to High Risk Youth* (U. of Minnesota: Minnesota Extension Service, 1986)
2. Gissler et. al., “Suicides After Pregnancy in Finland: 1987-94: register linkage study,” *British Medical Journal*, 313: 1431-1434, 1996; Campbell, et. al., “Abortion in Adolescence,” *Adolescence*, 23:813-823, 1988.
3. Gissler, op. cit.; Gissler, “Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000,” *European J. Public Health* 15(5):459-63, 2005. See also Reardon et. al., “Deaths Associated With Pregnancy Outcome: A Record Linkage Study of Low Income Women,” *Southern Medical Journal* 95(8):834-41, Aug. 2002.
4. Franz and Reardon, “Differential Impact of Abortion on adolescents and adults,” *Adolescence*, 27 (105), 172, 1992.
5. Somers, “Risk of Admission to Psychiatric Institutions Among Danish Women Who Experienced Induced Abortion: An Analysis Based on National Report Linkage” (Ph.D. Dissertation, Los Angeles: University of California, 1979, Dissertation Abstracts International, Public Health 2621-B, Order No. 7926066)
6. “Teenage Pregnancy: Overall Trends and State-by-State Information,” Report by the Alan Guttmacher Institute, Washington, DC, www.agi.org.
7. For more information, see www.unchoice.info.
8. Coleman, “Resolution of Unwanted Pregnancy During Adolescence Through Abortion Versus Childbirth: Individual and Family Predictors and Psychological Consequences,” *Journal of Youth and Adolescence* (2006).
9. Barglow and Weinstein, “Therapeutic Abortion During Adolescence: Psychiatric Observations,” *Journal of Youth and Adolescence*, 2(4):33, 1973.
10. Rue et. al., “Induced abortion and traumatic stress: A preliminary comparison of American and Russian women,” *Medical Science Monitor* 10(10): SR5-16, 2004.
11. Franz and Reardon, “Differential Impact of Abortion on adolescents and adults,” *Adolescence*, 27 (105), 172, 1992.
12. Belanger, et. al., “Pain of First Trimester Abortion: A Study of Psychosocial and Medical Predictors,” *Pain*, 36:339; Smith, et. al., “Pain of first-trimester abortion: Its quantification and relationships with other variables,” *American Journal Obstetrics & Gynecology*, 133:489, 1979.
13. Burkman, et. al., “Morbidity Risk Among Young Adolescents Undergoing Elective Abortion,” *Contraception*, 30(2):99, 1984; Schulz, et. al., “Measures to Prevent Cervical Injury During Suction Curettage Abortion,” *The Lancet*, 1182-1184, May 28, 1993.
14. Burkman, et. al., “Culture and treatment results in endometritis following elective abortion,” *American J. Obstet. & Gynecol.*, 128:556, 1997; Avonts and Piot, “Genital infections in women undergoing induced abortion,” *European J. Obstet. & Reproductive Biology*, 20:53, 1985; Cates, “Teenagers and Sexual Risk-Taking: The Best of Times and the Worst of Times,” *Journal of Adolescent Health*, 12:84, 1991.
15. “Teenage Pregnancy: Overall Trends and State-by-State Information,” Report by the Alan Guttmacher Institute, Washington, DC, www.agi.org.
16. Brind et. al., “Induced abortion as an independent risk factor for breast cancer: a comprehensive review and analysis,” *J. Epidemiology & Community Health*, 50:481, 1996.
17. Kochanck, “Induced Terminations of Pregnancy, Reporting States 1988,” *Monthly Vital Statistics Report*, 39(12): Suppl. 1-32, April 30, 1991.
18. Strahan, “Differential Adverse Impact on Teenagers Who Undergo Induced Abortion,” *Association for Interdisciplinary Research Bulletin*, 15(1):3, March/April 2000.
19. Strahan, “Psycho-Social Aspects of Late-Term Abortions,” *Assoc. For Interdisciplinary Research Bulletin*, 14(4):1, 2000.
20. Burkman, *American J. Obstet. & Gynecol.* op. cit; Lurie and Shoham, “Induced Midtrimester Abortion and Future Fertility: Where Are We Today?” *International J. of Fertility*, 40(6):311, 1995.
21. Atrash and Hogue, “The effect of pregnancy termination on future reproduction,” *Baillieres Clinic Obstet. & Gynecol.*, 4(2):391, 1990; Rooney, “Is Cerebral Palsy Ever a Choice?” *The Post-Abortion Review*, 8(4):4-5, Oct.-Dec. 2000.
22. Amaro, et al., “Drug use among adolescent mothers: profile of risk,” *Pediatrics*, 84, 1989, 144-150.
23. Horowitz, “Adolescent Mourning Reactions to Infant and Fetal Loss,” *Soc. Casework*, 59:551, 1978.
24. Smith, “A follow-up study of women who request abortion,” *American Journal of Orthopsychiatry*, 1973, 43: 574-585.
25. Zakus and Wilday, “Adolescent Abortion Option,” *Social Work in Health Care*, 12, 1987, 77-91.
26. Biro, et al., “Acute and Long-Term Consequences of Adolescents Who Choose Abortions,” *Pediatric Annals*, 15(10):667-672, 1986.
27. Campbell, et. al., “Abortion in Adolescence,” *Adolescence*, 23:813-823, 1988.
28. Wheeler, “Adolescent Pregnancy Loss,” in Woods, Jr. Woods (eds.), *Loss During Pregnancy or the Newborn Period* (1997); Cvejic et. al., “Follow-up of 50 adolescent girls 2 years after abortion,” *Canadian Medical Assoc. Journal*, 116:44, 1997.
29. Joyce, “The Social and Economic Correlates of Pregnancy Resolution Among Adolescents in New York by Race and Ethnicity: A Multivariate Analysis,” *American J. of Public Health*, 78(6):626, 1988.